
Contaminated Food Sickens 1 in 10 People Worldwide Each Year

04/12/2015



Giving its first global estimates of preventable foodborne diseases, a WHO report called on governments and industry to improve inspections and control of the food chain from the fields and farmyard to the factory and the plate.

Foodborne diseases - caused by bacteria such as salmonella, viruses, parasites, toxins and chemicals - mostly cause temporary symptoms such as nausea, diarrhea and vomiting.

But they can also cause longer-term illnesses including cancer, kidney or liver failure, brain disorders, epilepsy and arthritis, the United Nations agency said.

"The data we are publishing is only a very conservative estimate, we are sure that the real figure is bigger," Dr. Kazuaki Miyagishima, director of WHO's Department of Food Safety, told a news briefing.

Data curated by [HealthGrove](#)

Apart from estimating the extent of contamination, the report also looked at the way food is traded and sold.

See also:

- Mind: [Using Pigeons to Diagnosis Cancer](#) |
- Sustainability: [Plight of the Long-Distance Flyers](#) |
- Tech: [All Gene-Editing Research Should Proceed Cautiously, Scientists Conclude](#) |

- The Sciences: [New Law Paves the Way for Asteroid Mining--But Will It Work?](#)

Part of the problem can come from global trading in food, Miyagishima said: "If there is one country where food safety is weak and this country exports food to other countries, (it) becomes the weakest chain in the whole food production system."

Faulty handling at the other end of the food chain, for example by street vendors, is also a problem in many countries.

"It is much better to invest in training and education of street vendors than to try to penalize them," said University of Florida expert Dr. Arie Hendrik Havelaar. "That would be an important strategy for many countries in the world to improve the food safety situation."

The highest number of cases and deaths occur among the poor in developing countries, but the United States and Europe also have deadly outbreaks.

"Our results show that the biggest burden is in Africa and in southeast Asia, and there the death rates are highest, including those of children under five years of age," said Havelaar, who chaired the WHO group of 150 scientists that carried out the research for the report.

Children under age five account for 40% of the 600 million cases of foodborne disease worldwide a year and 125,000 of the 420,000 deaths, or 30%, the WHO said.

Children are especially vulnerable to diarrheal diseases, often caused by eating raw or undercooked meat, or eggs, fresh produce and dairy products that are contaminated, it said.

In Africa, most deaths are caused by salmonella, the pork tapeworm, cyanide in cassava and aflatoxin, a chemical produced by moulds that grow on improperly stored grains or corn.

Governments must invest more in training food producers, suppliers and the public, the WHO said.
